


Air-Sea Flux Impacts Over Misaligned Waves During Atmospheric Cold Fronts

César sauvage¹, Hyodae Seo^{1,2}, Benjamin W. Barr¹, James B. Edson¹ and Carol Anne Clayson¹

1-University of Hawai'i at Manoa, HI, USA

2-Woods Hole Oceanographic Institution



WOODS HOLE
OCEANOGRAPHIC
INSTITUTION



UNIVERSITY
of HAWAII
MĀNOA

WISE 2025 | Seattle, WA, USA

12-15 May 2025


Contact: csauvage@hawaii.edu



Introduction

Atmospheric cold fronts account for ~15% of the time during winter months in the North Atlantic Ocean. (Parfitt et al. 2017)

They are characterized by high wind speed, abrupt veering, and rapid translation.




Introduction

Atmospheric cold fronts account for ~15% of the time during winter months in the North Atlantic Ocean. (Parfitt et al. 2017)

They are characterized by high wind speed, abrupt veering, and rapid translation.

→ Leads to strongly misaligned waves behind the cold front




Introduction

Atmospheric cold fronts account for ~15% of the time during winter months in the North Atlantic Ocean. (Parfitt et al. 2017)

They are characterized by high wind speed, abrupt veering, and rapid translation.

→ Leads to strongly misaligned waves behind the cold front





What are the resulting impacts on the surface drag and the air-sea momentum flux?

Coupled Modeling and Observation

SCOAR Regional Coupled Modeling
(i.e., Seo et al. 2007, 2016, 2021 ; Sauvage et al. 2023)

WRF nested configuration: OUTER domain at 7.5 km
WRF-ROMS-WW3 COUPLED domain at 1.5 km





Coupled Modeling and Observation

SCOAR Regional Coupled Modeling
(i.e., Seo et al. 2007, 2016, 2021 ; Sauvage et al. 2023)

WRF nested configuration: OUTER domain at 7.5 km
WRF-ROMS-WW3 COUPLED domain at 1.5 km



Ocean Observatories Initiative
Pioneer Array (2014-2022)




Coupled Modeling and Observation

SCOAR Regional Coupled Modeling
(i.e., Seo et al. 2007, 2016, 2021 ; Sauvage et al. 2023)

WRF nested configuration: OUTER domain at 7.5 km
WRF-ROMS-WW3 COUPLED domain at 1.5 km



Ocean Observatories Initiative
Pioneer Array (2014-2022) 




National Data Buoy Center
(NDBC)




Atmospheric cold front passing on December 6th 2017

Aligned waves under the warm front

Strongly misaligned waves under the cold front



Atmospheric cold front passing on December 6th 2017




Waves stayed strongly misaligned more than 12h hours after the passage of the cold front ..!

Atmospheric cold front passing on December 6th 2017


Pre-cold-frontal

2017-12-06 09:00:00




Cold-frontal

2017-12-06 12:00:00



Post-cold-frontal


2017-12-06 15:00:00



10.0
1.0
0.1


NDBC

44066



Model

06-06:00 UTC



$E_f (\text{m}^2 \cdot \text{Hz}^{-1})$


10⁻³ 10⁻² 10⁻¹
Energy Density ($\text{m}^2 \cdot \text{s} / \text{deg}$)

■ Cold front location

Atmospheric cold front passing on December 6th 2017


Pre-cold-frontal

2017-12-06 09:00:00







Cold-frontal

2017-12-06 12:00:00



Post-cold-frontal

2017-12-06 15:00:00




■ Cold front location

NDBC


Model

44066

06-06:00 UTC



06-16:00 UTC



Momentum flux :

$$\tau = \rho u_*^2 = \rho C_D \Delta U^2$$

COARE 3.5 Wave based formulation (**WBF**) (Edson et al. 2013)

$$z_{rough} = H_s \cdot 0.09 \cdot \left(\frac{u_*}{C_p} \right)^2$$

→ Assumes waves aligned with winds

Momentum flux :

$$\tau = \rho u_*^2 = \rho C_D \Delta U^2$$

COARE 3.5 Wave based formulation (**WBF**) (Edson et al. 2013)

$$z_{rough} = H_s \cdot 0.09 \cdot \left(\frac{u_*}{C_p} \right)^2$$

→ Assumes waves aligned with winds



Modified COARE 3.5 Wave based formulation (**WBF** θ)
(Porchetta et al 2019, Sauvage et al. 2023)

$$z_{rough} = H_s \cdot 0.09 \cos(0.4\theta) \cdot \left(\frac{u_*}{C_p} \right)^{2 \cos(-0.32\theta)}$$

→ Takes into account wind and wave misalignment

Misaligned waves impact on momentum flux


2017-12-06-15:00



Surface drag coefficient and momentum stress increase by 20% and 11% on average behind the cold front

Misaligned waves impact on momentum flux


2017-12-06-15:00



Surface drag coefficient and momentum stress increase by 20% and 11% on average behind the cold front

Direct impact on the surface wind speed which decreases by 2% on average

Misaligned waves impact on heat fluxes



$$H_s = -\rho_a C_{pa} u_* T_*,$$


$$H_l = -\rho_a L_e u_* q_*,$$

$$R_r = \frac{u_* z_0}{\nu}, \quad z_{0t} = \frac{5.8e^{-5}}{R_r^{0.72}},$$

The scalar roughness Z_{0t} is decreased by 60%, leading to a decrease of T_* .

■ Cold front location

Misaligned waves impact on heat fluxes



$$H_s = -\rho_a C_{pa} u_* T_*,$$

$$H_l = -\rho_a L_e u_* q_*,$$


$$R_r = \frac{u_* z_0}{\nu}, \quad z_{0t} = \frac{5.8e^{-5}}{R_r^{0.72}},$$

The scalar roughness Z_{0t} is decreased by 60%, leading to a decrease of T_* .

The increase of u_* is of similar magnitude.

■ Cold front location

Misaligned waves impact on heat fluxes



$$H_s = -\rho_a C_{pa} u_* T_*,$$

$$H_l = -\rho_a L_e u_* q_*,$$

$$R_r = \frac{u_* z_0}{\nu}, \quad z_{0t} = \frac{5.8e^{-5}}{R_r^{0.72}},$$

The scalar roughness Z_{0t} is decreased by 60%, leading to a decrease of T_* .



The increase of u_* is of similar magnitude.

The overall effect on heat fluxes is generally small, reduced in average by 2~3%.

■ Cold front location

Direct covariance flux system measurements at Pioneer Array

20 atmospheric cold fronts detected during the period when momentum flux observations are available.



Using **WBF θ** generally increases the flux by 7% (up to 16%) in case of strongly misaligned waves after the cold front.

Taking into account misalignment generally reduces the bias from -7.6% in **WBF** to 3.5% in **WBF θ** .

Conclusion

- Atmospheric cold fronts lead to large areas of mixed sea conditions misaligned with local wind
- On average waves can stay misaligned for 5 to 10h after the passage of the cold front
- Surface drag and momentum are increased over the strongly misaligned waves
- Taking into account misalignment in the flux parameterization generally reduces the bias compared to observations

Conclusion

- Atmospheric cold fronts lead to large areas of mixed sea conditions misaligned with local wind
- On average waves can stay misaligned for 5 to 10h after the passage of the cold front
- Surface drag and momentum are increased over the strongly misaligned waves
- Taking into account misalignment in the flux parameterization generally reduces the bias compared to observations
- Keep working toward a better understanding of the misalignment in different wind waves regimes (i.e., high wind regimes, cross-swell vs opposed swell...)
- Need more long-term co-located observations (wind, waves, air-sea fluxes) like the Pioneer Array

Air-Sea Flux Impacts Over Misaligned Waves During Atmospheric Cold Fronts


César sauvage¹, Hyodae Seo^{1,2}, Benjamin W. Barr¹, James B. Edson¹ and Carol Anne Clayson¹

1-University of Hawai'i at Manoa, HI, USA

2-Woods Hole Oceanographic Institution



Sauvage et al. 2024



WOODS HOLE
OCEANOGRAPHIC
INSTITUTION



UNIVERSITY
of HAWAII
MĀNOA

WISE 2025 | Seattle, WA, USA

12-15 May 2025

Contact: csauvage@hawaii.edu



Supplementary Figures


Coupling frequency 1h
Horizontal resolution : 1.5 km
(7.5 km for OUTER domain)


Boundary forcing:

Atmospheric forcing:
hourly ERA5

Ocean forcing:
Daily Mercator Ocean Inter.

Wave forcing:
3-hourly spectral points
(Ifremer global runs)




Pre-cold-frontal**Post-cold-frontal**

Comparison of wave spectrum energy density at different NDBC mooring against our coupled model.

Strong southerly waves aligned with the wind in the pre-cold-frontal conditions.


While the wind shifted to northerly, strong southerly waves were still present leading to strong misalignment during the post-cold-front conditions.

To be noted that after the cold front, younger waves are generated following the northwesterly wind, contributing to the mixed sea state conditions.



Wave Directional Spectrum

Comparison of wave spectrum energy density at different NDBC moorings against our coupled model.




Strong southerly waves aligned with the wind in the pre-cold-frontal conditions.

While the wind shifted to northerly, strong southerly waves were still present leading to strong misalignment during the post-cold-front conditions.

To be noted that after the cold front, younger waves are generated following the northwesterly wind, contributing to the mixed sea state conditions.


Wave Spectrum Energy Density



Supplementary Figures

The surface exchange coefficients are increased by up to 5% behind the cold front.

The effect on heat fluxes is generally small, reduced in average by 2~3%.



$$\tau = \rho_a C_D \Delta U_r^2 = \rho_a u_*^2,$$


$$H_s = \rho_a C_{pa} C_h \Delta U_r \Delta T = -\rho_a C_{pa} u_* T_*,$$

$$H_l = \rho_a L_e C_e \Delta U_r \Delta Q = -\rho_a L_e u_* q_*,$$


$$R_r = \frac{u_* z_0}{\nu},$$

$$z_{0t} = \frac{5.8e^{-5}}{R_r^{0.72}},$$

Supplementary Figures



Supplementary Figures



Most of the cold fronts occur during the extended winter months, November-March

70% of the detected cold fronts lead to strongly misaligned waves.

Waves can stay misaligned between 5h to 10h after the passage of the cold front.
(depending on the chosen threshold to define alignment vs misalignment)

Supplementary Figures

